

2017-12-01

Pelagic habitat: exploring the concept of good environmental status

Dickey-Collas, M

<http://hdl.handle.net/10026.1/10101>

10.1093/icesjms/fsx158

ICES Journal of Marine Science

Oxford University Press (OUP)

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Pelagic habitat: exploring the concept of good environmental status.

Mark Dickey-Collas^{1,2*}, Abigail McQuatters-Gollop³, Eileen Bresnan⁴, Alexandra C. Kraberg⁵, John P. Manderson⁶, Richard D.M. Nash⁷, Saskia A. Otto⁸, Anne F. Sell⁹, Jacqueline F. Tweddle¹⁰, Verena M. Trenkel¹¹

¹ *International Council for the Exploration of the Sea, H.C. Andersens Boulevard 44-46, DK 1553, Copenhagen, Denmark*

² *DTU Aqua National Institute of Aquatic Resources, Technical University of Denmark (DTU), Kemitorvet 1, 2800 Kgs. Lyngby, Denmark, Denmark*

³ *Plymouth University, Centre for Marine Conservation and Policy, Drake Circus, Plymouth, Devon, PL4 8AA, UK*

⁴ *Marine Scotland Science, Marine Laboratory, 375 Victoria Road, Aberdeen, AB11 9DB, UK*

⁵ *Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung, Biologische Anstalt Helgoland, P.O. Box 180, 27483, Germany*

⁶ *NOAA/North East Fisheries Science Center Oceans & Climate and Cooperative Research Programs, 74 Magruder Road, Highlands, NJ 07732, USA*

⁷ *Institute of Marine Research, PB, 1870 Nordnes, 5817 Bergen, Norway*

⁸ *Institute for Hydrobiology and Fisheries Science, Center for Earth System Research and Sustainability, Hamburg University, Grosse Elbstrasse 133, 22767 Hamburg, Germany*

⁹ *Thünen Institute of Sea Fisheries, Palmallee 9, 22767 Hamburg, Germany*

¹⁰ *School of Biological Sciences, Zoology Building, University of Aberdeen, Tillydrone Ave, Aberdeen, AB24 2TZ, UK*

¹¹ *IFREMER, Rue de l'Île d'Yeu, BP 21105, 44311 Nantes, France*

*Corresponding author: e-mail: mark.dickey-collas@ices.dk

Keywords

Plankton, Marine Strategy Framework Directive, pelagic habitats, seascape, MPA

Abstract

Marine environmental legislation is increasingly expressing a need to consider the quality of pelagic habitats. This paper uses the European Union marine strategy framework to explore the concept of good environmental status (GES) of pelagic habitat with the aim to build a wider understanding of the issue. Pelagic ecosystems have static, persistent and ephemeral features, with manageable human activities primarily impacting the persistent features. The paper explores defining the meaning of “good”, setting boundaries to assess pelagic habitat and the challenges of considering habitat biodiversity in a moving medium. It concludes that for pelagic habitats to be in GES and able to provide goods and services to humans, three conditions should be met: i) all species present under current environmental conditions should be able to find the pelagic habitats essential to close their life cycles; ii) biogeochemical regulation is maintained at normal levels; iii) critical physical dynamics and movements of biota and water masses at multiple scales are not obstructed. Reference points for acceptable levels of each condition and how these may change over time in line with prevailing oceanographic conditions, should be discussed by knowledge brokers, managers and stakeholders. Managers should think about a habitat hydrography rather than a habitat geography. Setting the bounds of the habitats requires a consideration of dimension, scale and gradients. It is likely that to deal with the challenges caused by a dynamic environment and the relevance of differing spatial and temporal scales, we will need to integrate multidisciplinary empirical data sets with spatial and temporal models to assess and monitor progress towards, or displacement from GES of the pelagic habitat.

45 Introduction

46 In 2008, the European Union enacted a novel piece of legislation requiring its countries to define, and then
47 monitor progress towards achieving, good environmental status (GES) for, amongst other things, pelagic
48 habitats (European Commission, 2008, 2010). This legislation is called the Marine Strategy Framework
49 Directive (MSFD, Bigagli *et al.*, 2015) and it sits within a patchwork of European legislation designed to
50 protect and encourage sustainable exploitation of the marine environment under the Integrated Maritime
51 Policy (Apitz *et al.*, 2006; European Commission, 2007; Boyes and Elliot, 2014). The Directive provides a
52 framework of guidance and actions for EU member states. For a range of anthropogenic pressures, and
53 states of the marine environment, each country is asked to define “good environmental status” as a target
54 and make the binary decision of whether they have achieved it or not (Borja *et al.*, 2010). If the answer is
55 that they have not, the countries should implement management measures to ensure that they will reach
56 GES.

57 Under descriptor 1 of the MSFD, which covers biodiversity, countries have to consider GES of habitats. This is
58 not dissimilar to Essential Fish Habitat (EFH) in USA legislation. Under the provisions of the Magnuson
59 Stevens Fisheries Conservation and Management Act (NOAA, 1996), a statutory mandate requires all
60 fisheries management plans to include descriptions of “essential fish habitat”, to identify adverse fishing
61 impacts and to conserve and enhance EFH. “Essential fish habitat” is defined as waters and substrate
62 necessary to fish for spawning, breeding, feeding or growth to maturity. In 2017, a new EU decision was
63 published (European Commission, 2017) which further described what should be considered around the
64 quality of pelagic habitat under the MSFD. When compared to other components of the marine ecosystem,
65 the impact of human activities on pelagic systems may appear minimal (Papathanasopoulou *et al.*, 2016),
66 but no consensus exists regarding the definition of GES for pelagic habitats. A clear understanding of the
67 attributes required for pelagic habitats to be in GES is required to guide monitoring and management
68 objectives.

69 This paper will use the arena of the MSFD to explore what is good environmental status for pelagic habitat,
70 and we hope that this example will provide useful insights for other similar legislative higher order objectives
71 for pelagic habitat, such as EFH and the like.

72 The challenge and the legislation

73 Knowledge brokers are being asked to provide guidance on what is an ecosystem in a good or bad state, a
74 question that is intrinsically normative (Turnhout *et al.*, 2007). The phrase “good environmental status”
75 means different things to different people and is value laden. It is probable that a decade ago, we would
76 have been discussing stewardship of “productive pelagic ecosystems”, but the MSFD uses concepts which
77 require public support and is also prone to moving social norms (Mee *et al.*, 2008). Tett *et al.* (2013) draw
78 parallels between the use of GES and the phrase “ocean health”, suggesting that the terms are metaphors
79 for a vision that aggregates over system components. The MSFD and various studies attempt to aid the
80 decision and assessment process by providing descriptive guidance (Mee *et al.*, 2008; Borja *et al.*, 2013; Tett
81 *et al.*, 2013). The guidance for habitats in the MSFD states

82 *“Biological diversity is maintained. The quality and occurrence of habitats and the distribution and*
83 *abundance of species are in line with prevailing physiographic, geographic and climatic conditions.”*
84 (European Commission, 2008).

85 This is further expanded in the 2010 supporting decision

86 *“the term habitat addresses both the abiotic characteristics and the associated biological community,*
87 *treating both elements together in the sense of the term biotope.... The three criteria for the assessment of*

88 *habitats are their distribution, extent and condition (for the latter, in particular the condition of typical*
89 *species and communities), accompanied with the indicators related respectively to them.”* (European
90 Commission, 2010).

91 The 2017 revised decision says that the condition of “*Pelagic broad habitat types (variable salinity, coastal,*
92 *shelf and oceanic/beyond shelf) ... including biotic and abiotic structure and functions ... is not adversely*
93 *affected due to anthropogenic pressures”* (European Commission, 2017). This should be assessed at the scale
94 of habitat adversely affected in square kilometres (km²) and as a proportion (percentage) of the total extent
95 of the habitat type. In the term GES, the word “good” is in relation to humans and thus linked to the
96 provision of goods and services, and stewardship and conservation for future generations (intergenerational
97 equity). The 2017 revision introduces the concept of habitat adversely affected by anthropogenic pressures.
98 In addition, any definition of good or adversely affected is often influenced by the suite of data, readily
99 available, with which to produce metrics as indicators of a pelagic habitat rather than an operational
100 definition of GES.

101 Similar to “good”, what “essential” means could also be considered normative. The Magnuson Stevens
102 Fisheries Conservation and Management Act emphasizes the quality of habitats with respect to effect on
103 growth, reproduction, and/or survival of different life stages and ultimately on the productivity of fishery
104 species. The definition of EFH is therefore organism-centered rather than anthropocentrically defined, and
105 integrates both pelagic and benthic habitats.

106 Diverse services are provided by the marine pelagic habitat (or combined habitats) such as the regulation of
107 ocean circulation and weather, carbon recycling and balance, production of living resources, and tourism.
108 Any consideration of good pelagic habitat needs to relate to the perceived priorities and objectives of
109 society. Any consideration of adverse pressures, needs to be in relation to some framework.
110 Anthropocentric, societal definitions of marine habitat and habitat quality can lead to the misclassification of
111 marine habitats based upon terrestrial analogies and teleologies. The possibility of falling into traps of
112 misclassification is particularly high in pelagic ecosystems that are embedded in turbulent heterogeneous
113 liquids.

114 **Igniting the discussion**

115 An open theme session was held at the 2016 ICES annual science conference in Riga, Latvia with the title
116 “What is a good pelagic habitat?” ([http://ices.dk/news-and-events/asc/ASC2016/Pages/Theme-session-](http://ices.dk/news-and-events/asc/ASC2016/Pages/Theme-session-J.aspx)
117 [J.aspx](http://ices.dk/news-and-events/asc/ASC2016/Pages/Theme-session-J.aspx)). The session was advertised to address the demands for clearer understanding on what is good
118 pelagic habitat as society is asking for guidance on what is a good or bad pelagic system. The focus of the
119 session was on the higher order objectives and was attended by 20-40 participants. Various presentations
120 illustrated the services provided by pelagic habitats such as the regulation of ocean circulation and weather,
121 carbon recycling and balance, production of living resources, importance of species or functional biodiversity
122 and tourism. The session split into three subgroups to consider what is meant by good pelagic habitat, how
123 can we quantify it and what are the features that distinguish it from other habitats? At the end of the
124 session, all participants were invited to contribute to the construction of a food for thought article building
125 on the ideas discussed. This paper is the result of the process.

126 **Pelagic habitat**

127 Pelagic habitats, following the MSFD definition as a biotope used in this paper, can be viewed as having
128 faster dynamics and lower levels of predictability when compared to terrestrial and marine benthic habitats
129 (Ray, 1991; Gray, 1997; Hyrenbach *et al.*, 2000). Living in liquid is different from living in gas, and the vital
130 rates of all marine organisms are controlled to a great degree by the properties of and processes occurring
131 within the ocean’s turbulent liquid (Purcell, 1977; Andersen *et al.*, 2015; Manderson, 2016). Bertrand et al.

(2014) describe the pelagic ecosystem as that where the 'substrate' consists of constantly moving water masses, where ocean surface turbulence creates ephemeral oases. Pelagic habitats are also defined by the frontal structures and subsides created and delivered by divergent and convergent flows (Tew Kai, 2009; Della Penna, 2017). The combination of the properties of pelagic systems has led to a formalisation of 'seascape ecology' as opposed to, the rather different, terrestrial 'landscape ecology' (Manderson, 2016). Ban et al (2014) highlight for the ocean system, that many species are widely distributed and wide ranging; the sizes and boundaries of biogeographical domains vary significantly by depth; habitat types exhibit a range of stabilities, from ephemeral (e.g., surface frontal systems) to hyper-stable (e.g. deep sea); and vertical and horizontal linkages are prevalent.

It could be said that a holistic approach that does not compartmentalise habitats (i.e. not treating benthic and pelagic habitat separately) is more in keeping with an integrated management. Within and across life history stages, many marine organisms are obligate integrators of benthic and pelagic properties and processes occurring within the oceans. However, most biodiversity legislation (including the MSFD) requires habitats to be defined and delineated on the basis of a patch-based view of seascapes analogous to the operational paradigm of terrestrial landscape ecology (Ray, 1991). This aids assessment, targeted management action, and communication of relevant issues. There are techniques to define boundaries between pelagic habitats using hydrographic variables and their spatial gradients calculated at an appropriate spatial scale (see Alvarez-Berastegui *et al.*, 2014). It is useful to think about pelagic habitats in terms of the static, persistent and ephemeral aspects (Hyrenbach *et al.*, 2000), with the static being bathymetric and coastal features, the persistent being hydrographic and climatic features which often vary seasonally, and the ephemeral being short lived and less predictable gradients in water qualities (Hyrenbach *et al.*, 2000). Other classification approaches to pelagic habitats exist (see Kavanaugh *et al.*, 2016) and the concept of gradient approaches is beginning to be considered even in terrestrial ecology (McGarigal and Cushman 2005; Cushman *et al.*, 2010) but we chose to keep the classification simple. Biological features can similarly be considered across these environmental axes of the seascape (Hidalgo *et al.*, 2015).

The MSFD mentions "prevailing physiographic, geographic and climatic conditions" (European Commission, 2008). In pelagic habitats, these prevailing conditions can be highly variable and dominate our observations of trends in state (McQuatters-Gollop, 2012). Pelagic organisms are embedded in a turbulent advective environment; their size determines how they are affected by the properties of the liquid and their scales of variability (e.g. effect of Reynolds number, advection or migration, etc., Kavanaugh *et al.*, 2016). Their behaviour and self-organisation (e.g. schooling behaviour) also impact their distribution in relation to physical/environmental forcing (see figure 10 in Bertrand et al 2008). Hidalgo *et al.* (2015) suggest that the effect of static and ephemeral features on our observations of biodiversity is often overridden by different non-linear effects in the pelagic environment. Predictability is challenged by the dynamics of the system. In most pelagic systems the prevailing conditions are a consequence of bathymetry, location, relative depth, temperature, salinity, oxygen, circulation, ice cover, carbon dioxide, light and turbidity. Many of these properties are highly dynamic because they are strongly forced directly or indirectly by the dynamics of the atmosphere and planetary motions. The consequences of the behaviour of organisms and the issue of scale (temporal and spatial) further complicate any assessment of habitat quality (e.g. Bertrand *et al.*, 2010; Louzao *et al.*, 2011; Miller *et al.*, 2015; Cisewski and Strass, 2016).

Many species inhabit the water column only temporarily such as meroplankton, mysids, or benthopelagic fish. Other species migrate over long distances or between coastal and offshore areas at daily to multi-annual time scales. Consequently, understanding the composition and trophic structure shared by a set of interacting communities and its dynamical implications for the persistence of biodiversity remains challenging (Melián *et al.*, 2005).

Scale, monitoring and boundaries

178 When monitoring and assessing the pelagic ecosystem care needs to be taken about the relevant scales,
 179 both spatial and temporal (see figure 3 in Kavanaugh *et al.*, 2016). The concept of scale was recently
 180 highlighted as one of the most useful ecological concepts to emerge in the last 100 years of ecological
 181 research (Reiners *et al.*, 2017) but it is also one of the most challenging when applying ecological concepts
 182 into operational management (see Stommell, 1963; Steele, 1978; Ban *et al.*, 2014). Temporally, the pelagic
 183 ecosystem varies within a day (e.g. diel migration, tidally driven changes in turbulence), across seasons (e.g.
 184 stratification), years and even multi-decadal cycles too. These cycles impact the persistent and ephemeral
 185 features. Spatially, variation of communities can range across many scales (Scales *et al.*, 2017). Since
 186 Schneider (2001) suggested that little is known about the importance of small- and large-scale processes on
 187 the structure of communities, progress has been made in understanding the dynamics and distribution of
 188 pelagic organisms across their habitat (Alvarez-Berastegui *et al.*, 2014; Bertrand *et al.*, 2014; Scales *et al.*,
 189 2017).

190 Tett *et al.* (2013) define good ecosystem status (good ocean health) as “*the condition of a system that is self-
 191 maintaining, vigorous, resilient to externally imposed pressures, and able to sustain services to humans. It
 192 contains healthy organisms and populations, and adequate functional diversity and functional response
 193 diversity. All expected trophic levels are present and well interconnected, and there is good spatial
 194 connectivity amongst subsystems.*” This definition requires an understanding of open marine systems and
 195 the interconnections between static elements and sub-systems. Pelagic habitats usually do not have distinct
 196 boundaries and are often defined by latitudinal and hydrographic gradients, semipermeable frontal
 197 boundaries between different water masses, and defined differences in density and current flows which may
 198 be seasonally variable (Alvarez-Berastegui *et al.*, 2014; Hidalgo *et al.*, 2015). Inshore, the relevant dynamics
 199 can be constrained by the geometry and geography of coastlines and the seabed along with characteristic
 200 seasonal frequencies of frontal formation and disintegration and associated changes in temperature,
 201 precipitation, and winds as well as tidal forcing. Concepts of GES, therefore, need to be spatially and
 202 temporally relevant to specific ecological processes or ecosystem services (Mee *et al.*, 2008). As with
 203 integrated ecosystem assessments, setting of boundaries is a key stage of an assessment of habitat (Dickey-
 204 Collas, 2014). The ideas behind conservation of habitat diversity and the role of functional redundancy in
 205 maintaining ecosystem resilience have been heavily influenced by research performed in terrestrial systems,
 206 shallow water reefs and benthic communities. These properties of promoting resilience are probably equally
 207 important in pelagic systems but less easily defined. Gray (1997) emphasised that it was important to
 208 consider the issue of scale in seascape diversity as relevant scales are determined by the specific ecological
 209 or ecosystem process. The spatial aspects of the MSFD (the subregions, the lack of coverage in the high seas
 210 and limits in coastal waters) may not be robust enough to cope with the range of spatial scales of pelagic
 211 habitat (see Ban *et al.*, 2014). The results of analysing temporal trends can be affected by the spatial scale of
 212 monitoring, with incorrect assessments if linear relationships are assumed (Bartolino *et al.*, 2012). The
 213 metrics used as indicators to assess and monitor GES have yet to be sufficiently tested for their robustness
 214 and applicability at different spatial scales (e.g. Wasmund *et al.*, 2017). Current monitoring of pelagic
 215 ecosystems generally does not exist at the spatial or temporally relevant scales necessary to assess
 216 prevailing conditions and some fine or large scale anthropogenic pressures. However, rapid advances in
 217 technology and the implementation of various levels of ‘Ocean Observation Systems’ are making the
 218 attainment of appropriate observational and monitoring data achievable (e.g. Kavanaugh *et al.*, 2016;
 219 Manderson, 2016; Trenkel *et al.*, 2016). Further advances are being made to develop the monitoring and
 220 statistical methods to assess the interaction of scale and habitat (Mayor *et al.*, 2007; Pittman and Brown,
 221 2011).

222 With increasing accuracy, we can model the impact of global, regional and local events in the pelagic system
 223 and explore future scenarios in relation to prevailing conditions and changes in pressures (Fernandes *et al.*,
 224 2013; Akimova *et al.*, 2016; Queirós *et al.*, 2016). Hufnagl *et al.* (in press) investigated 10 different physical

models for the oceanography of the southern North Sea and suggested that most models showed systematic biases during all years in comparison to the ensemble median, indicating that, in general, inter-annual variation was represented equally by the models but absolute values of movement and temperature experienced by particles varied when modelling particles through the system. We can also determine aspects of connectivity with an appropriate spatial scale of dispersal, and the broad scale influence of oceanography on near shore oceanographic dispersal variability (Watson *et al.*, 2011; Treml *et al.*, 2012). Monitoring, assessment and the setting of thresholds need to be designed/accountable for this variability, probably by using targeted finer scale monitoring of areas of concern within broader scale seascape integrated modelling of larger regions.

Assessments for management

Even with a definition of GES, the variability in prevailing conditions makes reaching the GES target challenging. It is also often unclear which human activities are putting pressure on the state of the pelagic ecosystem (Shephard *et al.*, 2015). The revised MSFD decision (European Commission 2017) states that the pelagic habitat must not be adversely affected due to anthropogenic pressures. When considering the pelagic habitat, it is important to consider the influence of upstream events. When assessing GES and where we are in relation to it, many researchers propose the use of the Driver-Pressure-State-Impact-Response (DPSIR) framework to guide management measures (Gimpel *et al.*, 2013; Knights *et al.*, 2013). This assumes that there are direct levers that can be pulled to reduce or increase the human pressures resulting in ecosystem response in a predicted direction. This poses problems when prevailing conditions are thought to have more impact on the pelagic system than any direct consequence of a human-caused pressure (McQuatters-Gollop, 2012). The obvious example of a clear DPSIR relationship is how fishing and hunting influence populations and ecosystem structure (Shephard *et al.*, 2014). However, interactions with other drivers often complicate such a clear relationship, making causal relationships more difficult to disentangle, e.g. in the case of fishing pressure and climate change acting simultaneously (Planque *et al.*, 2010; Planque, 2015). However, when the influence of anthropogenic pressures is less easy to detect, such as when prevailing conditions play a strong role in habitat dynamics, surveillance indicators can be used to monitor pelagic community structure (Shephard *et al.*, 2015). If a surveillance indicator shows an unwelcome trajectory, beyond predefined thresholds, management action should be triggered. But the lack of defined GES for pelagic habitat means that the objectives for monitoring and action are not so clear.

Good environmental status of the pelagic habitat is not synonymous with setting up a marine protected area (MPA) for pelagic habitat. The latter can be seen as a tool to help achieve the former (Game *et al.*, 2009). Pelagic MPAs have tended to focus on biodiversity or productivity “hotspots” (Etnoyer *et al.*, 2004; Scales *et al.*, 2014). The behaviour of animals is often explored in relation to oceanography and geography (e.g. Vilchris *et al.*, 2006; Kobayashi *et al.*, 2008; Louzao *et al.*, 2011) providing information of relevant areas in need of protection. Ban *et al.* (2014) explore this further (see Figure 1 in their paper). The MSFD clearly states that there should be no further loss of diversity in genes, species or habitats (Borja *et al.*, 2013), and goods and services are also derived from pelagic habitats not associated with biodiversity hotspots. It is therefore as important to conserve the low biodiversity habitats as the hotspots (Gray, 1997, e.g. the central Arctic Ocean, and estuaries), requiring a toolset wider than MPAs alone.

A pelagic habitat can also be in a natural ecological state even when that state may be perceived to be ‘negative’ by societies. In some areas, the accumulation of high concentrations of algal toxins in shellfish can be driven by natural forces (prevailing conditions) but considered by society as ‘negative’ owing to the economic impact resulting from enforced closures of shellfish harvesting areas (Gowen *et al.*, 2012). Similarly, high biomass blooms of the dinoflagellate *Karenia mikimotoi* can result in mortalities of the benthos or farmed fish; however, these events may be driven by natural bloom formation offshore and transport in coastal currents (Davidson *et al.*, 2009; Gillibrand *et al.*, 2016) and not human activities. Because

271 marine ecosystems are nonlinear with complex feedback loops and multiple stable states, cyclic disturbances
272 may cause collapses in ecosystem states due to changes in natural oceanographic forcing. Such collapses can
273 be perceived to be negative by humans in the short term. However they may in fact be necessary for
274 periodically resetting some ecosystem trajectories toward “healthier” states. Oceanographic disturbance
275 and ecosystem state collapses related to El Niño and La Niña cycles are hypothesized to underlie ecosystem
276 dynamics in the highly productive Peruvian upwelling system (Bakun and Weeks, 2008). Ecosystem dynamics
277 resulting from prevailing oceanographic conditions need to be distinguished from those resulting from
278 human impacts, particularly eutrophication (Gowen *et al.*, 2012) and pollution events (e.g. oil spills).
279 Although defining good environmental status is normative, when setting targets we must avoid labelling
280 natural but unwished for conditions as Bad Environmental Status.

281 **What is good pelagic habitat?**

282 The contributions to the open theme session and the subgroup discussions described above provided the
283 input material for considering the requirements for GES for pelagic habitats. The issues discussed included
284 retaining sustainable exploitation and a resilient ecosystem. A comprehensive definition of a resilient
285 ecosystem remains elusive, however, here we consider resilience as an ability of the ecosystem to return to
286 a state from which it was perturbed. The aim of the exploration was to find a pragmatic approach to ensure
287 resilience and sustainability using tangible and operational phraseology. The key services offered and
288 properties required from pelagic habitats were then considered and selected based on expert knowledge
289 and information in the literature. The identified services related to regulation and habitat functions as
290 defined for example by de Groot *et al.* (2002). They included services provided by all habitats, terrestrial and
291 marine, as well as services more specific to the pelagic habitat.

292 Life cycle maintenance is considered an essential marine and coastal ecosystem service (Liquete *et al.* 2013).
293 For considering this habitat function the framework developed by Petitgas *et al.* (2013) to analyse climate
294 impacts on habitats was viewed as useful, in that it allows for an analysis of habitat requirement by life
295 stage. Because it provides linkages between and integrates requirements across life stages, such a
296 framework could be developed for assessing the status of pelagic habitats. In addition, it can be used to
297 assess impacts across the entire life cycle, including where necessary information on benthic-pelagic
298 connectivity through organisms that use the both benthic and pelagic habitats at different life stages. This
299 would be applicable when considering both exploitation and conservation objectives.

300 Pelagic habitats contribute to the functioning of the global bio-geochemical system, in particular to ocean
301 nourishment (Liquete *et al.* 2013). The main services are nutrient cycling (for example C, O, N, P, S, Si, Fe)
302 and gas regulation (Costanza *et al.* 1997). The oceans have been a net sink of increased atmospheric carbon
303 dioxide, with ocean warming expected to reduce this role (see chapter 13 Millenium Ecosystem Assessment
304 2005 <http://www.millenniumassessment.org/documents/document.282.aspx.pdf>). The sea surface – air
305 interface, the upper boundary of the pelagic habitat, plays an important role in this gas exchange between
306 the ocean and the atmosphere. Within the pelagic habitat, growing and moving organisms contribute to
307 nutrient transportation and recycling. Algal blooms will occur naturally, however, in coastal waters along
308 with hypoxia are signs of pollution and eutrophication surpassing system capacity and an immediate
309 resilience for suppressing catastrophic events. Linkages between marine and terrestrial ecosystems occur
310 because the major human activities impacting these services are land based (MEA 2005).

311 Linked to both of these functions is the inherent physical nature of the liquid substrate. The physical
312 qualities of the pelagic habitat warrant additional attention, including temperature, salinity and energy
313 gradients. As highlighted by Ban *et al.* (2014), Hidalgo *et al.* (2015) and Scales *et al.* (2017) the pelagic habitat
314 is structurally different from terrestrial and benthic, i.e. solid habitats. The unique properties and the
315 consequences of the pelagic habitat needs to be incorporated into any assessment of GES. These properties
316 contribute to the wider habitat function.

Following on from these considerations, we offer a very simple concept when defining GES for pelagic ecosystems. It is possible to consider the pelagic habitat as hydrography-driven, rather than geography-driven. This means that specific conditions and habitats are not fixed in space or time. This concept will allow scoping for national/regional definitions of GES. We are aware that the MSFD expects future anthropogenic impact and economic and social development of the seas; the MSFD does not strive towards returning European marine waters to a pristine state. What actually constitutes a pristine state is a matter of much debate due to the long and short term dynamic nature of the environment. Instead, the MSFD emphasises sustainable use of the marine system and recognises that GES should be a realistic and attainable target (European Commission, 2008). This contrasts to the organism focused Essential Fish Habitat concept. Here we suggest three key overlapping and interactive properties of the state of the system that ensure essential services that must be prioritised as contributors to GES.

1. Life cycle closure for marine organisms.

Central to the provision of goods and services, and conservation priorities, is that pelagic habitats maintain their ability to act as reproduction (including spawning and mating), nursery, and feeding grounds, as well as migration and advection routes, for marine organisms, resulting in no further decrease in global and regional natural biodiversity in line with prevailing oceanographic conditions. This includes organisms that spend all life stages in the water column and those that use it during various stages of their life cycle. For generational equity, no species - with its essential habitat - should be threatened by anthropogenic activity. This property produces what is called a habitat ecosystem service (e.g. Costanza *et al.* 1999, de Groot *et al.* 2002).

2. The global and regional roles of pelagic systems in biogeochemical regulation.

The pelagic ecosystem fulfils a wide range of roles in the regulation, recycling, transfer, storage and release of biochemical components and processes of relevance to global, regional and localised health of the seas and the whole planet. These roles include cycling of carbon, oxygen, nutrients, carbon sequestration, and many others. When determining GES, these roles must be acknowledged. The biochemical functions of the pelagic system should not move beyond what is considered normal under prevailing climatic conditions, supporting the key structural and functional aspects of pelagic ecosystems.

3. The physically dynamic nature of pelagic habitats.

The pelagic system provides movement of energy and materials that are important at the global, regional and local scales. GES should account for this role of the liquid in determining trophic and life cycle coupling. The movement of water, the interaction with weather, the provision of renewable energy, the advection of substances, coastal erosion, etc., are all relevant to the definition of GES. Consideration of pelagic habitat state must consider both Lagrangian and Eulerian aspects of that state, thus an awareness of the impact of upstream and consequences for downstream events. Whilst most of this movement and impacts of hydrodynamics is not manageable at anything except the local scale, a recognition that movement of organisms, materials and energy is a key part of pelagic habitats must be included in GES considerations.

For all of the three to be achieved all anthropogenic activities and pressures need to be managed or mitigated and the influence of physics understood (Ban *et al.*, 2014). This management includes achieving or maintaining low anthropogenic nutrient input maintaining stoichiometry of elements and minimizing the introduction of litter (including plastic), near zero contaminant pollution, and sustainable fishing; maintaining healthy plankton communities; and due consideration for siting permanent marine structures and regulating marine traffic, to maintain efforts to reduce introduced non-native and invasive species

(OSPAR, 2010; HELCOM, 2010). This requires management measures for the terrestrial landscape to be enacted too, as pressures are often sourced up stream on land.

Salience, legitimacy and credibility

This food for thought article was written by scientists with an interest in research in the pelagic ecosystem. Some of us work closely at the science/policy interface. We wrote this paper to stimulate discussion about higher order objectives for the pelagic habitat (Jennings, 2005), and it can be seen as an initiation of a dialogue between scientists and society (recognised as Mode 2 science by Gibbons *et al.*, 1994). Under the MSFD, the definition of GES is the responsibility of EU member states, hopefully working together through the European Regional Seas conventions (e.g. OSPAR and HELCOM). We would hope that any setting of a vision for pelagic GES would involve a scoping process between knowledge brokers, managers, and stakeholders. A similar exercise took place to explore the ecosystem approach objectives for pelagic fisheries (Trenkel *et al.*, 2015), where two independent scoping exercises gave remarkably similar results for potential higher order objectives.

However, in contrast to the exploration of higher order objectives carried out by Trenkel *et al.* (2015), we did not scope with stakeholders from beyond the scientific realm and limited the exploration to scientists joining the dedicated theme session by interest, without attempting to balance expertise of attendees. This leads us to likely criticism in terms of the salience, legitimacy and credibility of our message (Cash *et al.*, 2002). The word 'salience' requires that the intervention by a group is appropriate at the time, and we argue that the MSFD being executed in Europe makes such a discussion very relevant. However, we acknowledge that our intervention lacks much legitimacy, because we have not engaged in wider stakeholder dialogue and we do not formally represent society as self-appointed interested parties. As scientists with an interest in pelagic research, and an interest in applied research, it is valid to question our motives to initiate the discussion. We have an interest in the profile of the issue being raised, i.e. we are clearly stakeholders (Funtowicz and Ravetz, 1993). By using a session at the ICES annual science conference, we have attempted to create an open arena for the discussion amongst the scientific community. We have sought to improve our credibility by describing our methods and publishing this article in a peer reviewed journal. The idea behind the article was to provide a resource to enable discussion with a broader stakeholder community.

Conclusion

Pelagic ecosystems have static, persistent and ephemeral features, with manageable human activities impacting, primarily, persistent features. Managers should think about a habitat hydrography rather than a habitat geography. Setting the bounds of the habitats requires a consideration of dimension, scale and gradients. For pelagic habitats to be in GES and able to provide goods and services to humans, three conditions should be met for pelagic waters: i) all species present under current environmental conditions have access to the pelagic habitats essential to close their life cycles; ii) biogeochemical regulation is maintained at normal levels; iii) critical physical dynamics and movements of biota and water masses at multiple scales are not obstructed. Reference points for acceptable levels of each condition and how these may change over time in line with prevailing oceanographic conditions, need to be discussed by knowledge brokers, managers and stakeholders. It is likely that to deal with the challenges caused by a dynamic environment and the relevance of differing spatial and temporal scales, we will need to integrate multidisciplinary empirical data sets with spatial and temporal models to assess and monitor progress towards, or movement from, GES of the pelagic habitat.

Acknowledgements

This food for thought contribution was initiated by theme session J "What is a good pelagic habitat?" at the 2016 ICES annual science conference <http://www.ices.dk/news-and-events/asc/ASC2016/Pages/Theme->

405 [session-J.aspx](#). All who took part in the session are thanked for their contributions. All participants were
406 welcome to contribute to this article. J. F. Tweddle was supported by the Natural Environment Research
407 Council [NERC grant reference number NE/P005756/1]. E. Bresnan was supported by the Scottish
408 Government's schedules of service ST02a and ST03r. A. McQuatters-Gollop was support by the Natural
409 Environment Research Council [NERC grant reference number NE/L002663/1].

410 References

- 411 Alvarez-Berastegui, D., Ciannelli, L., Aparicio-Gonzalez, A., Reglero, P., Hidalgo, M., López-Jurado, J. L.,
412 Tintoretto, J. And Alemany, F. 2014. Spatial scale, means and gradients of hydrographic variables define
413 pelagic seascapes of bluefin and bullet tuna spawning distribution. PLoS one, 9(10), e109338. 14pp
- 414 Akimova, A., Núñez-Riboni, I., Kempf, A., and Taylor, M. H. 2016. Spatially-resolved influence of temperature
415 and salinity on stock and recruitment variability of commercially important fishes in the North Sea. PLoS
416 ONE, 11(9): e0161917. doi:10.1371/ journal.pone.0161917.
- 417 Andersen, A., Wadhwa, N., and Kiørboe, T. 2015. Quiet swimming at low Reynolds number, Physical Review
418 E, 91, 042712.
- 419 Apitz, S. E., Elliott, M., Fountain, M., and Galloway, T. S. 2006. European Environmental Management:
420 Moving to an Ecosystem Approach. Integrated Environmental Assessment and Management, 2(1): 80-85.
- 421 Bakun, A., and Weeks, S.J., 2008. The marine ecosystem off Peru: What are the secrets of its fishery
422 productivity and what might its future hold? Progress In Oceanography 79:290–299.
- 423 Ban, N. C., Maxwell, .S.M., Dunn, D.C., Hobday, A.J., Bax, N., Ardron, J., Gjerde, K.M., Game, E.T., Devillers, R.,
424 Kaplan, D.M., Dunstan, P.K., Halpin P.N., Pressey, R.L. 2014. Better integration of sectoral planning and
425 management approaches for the interlinked ecology of the open oceans. Marine Policy 49: 127-136
- 426 Bartolino, V., Ciannelli, L., Spencer, P., Wilderbuer, T. K., & Chan, K. S. 2012. Scale-dependent detection of
427 the effects of harvesting a marine fish population. Marine Ecology Progress Series, 444, 251-261.
- 428 Bertrand, A., Ballon, M., and Chaigneau, A. 2010. Acoustic observation of living organisms reveals the upper
429 limit of the oxygen minimum zone. PLoS ONE, 5(4): e10330. doi:10.1371/journal.pone.0010330.
- 430 Bertrand, A., Gerlotto, F., Bertrand, S., Gutiérrez, M., Alza, L., Chipollini, A., Díaz, E., Espinoza, P., Ledesma, J.,
431 Quesquén, R., Peraltila, S., Chavez, F. 2008. Schooling behaviour and environmental forcing in relation to
432 anchoveta distribution: An analysis across multiple spatial scales. Progress in Oceanography 79: 264–277.
- 433 Bertrand, A., Grados, D., Colas, F., Bertrand, S., Capet, X., Chaigneau, A., ... Fablet, R. 2014. Broad impacts of
434 fine-scale dynamics on seascape structure from zooplankton to seabirds. Nature communications, 5.
435 Article Number: 5239.
- 436 Bigagli, E. 2015. The EU legal framework for the management of marine complex social–ecological systems.
437 Marine Policy, 54: 44–51.
- 438 Borja, A., Elliott, M., Carstensen, J., Stiina Heiskanen, A., and van de Bund, W. 2010. Marine management –
439 Towards an integrated implementation of the European Marine Strategy Framework and the Water
440 Framework Directives. Marine Pollution Bulletin, 60: 2175–2186.
- 441 Borja, A., Elliott, M., Andersen, J. H., Cardoso, A. C., Carstensen, J., Ferreira J. G., Heiskanen, A. S., *et al.* 2013.
442 Good Environmental Status of marine ecosystems: What is it and how do we know when we have
443 attained it? Marine Pollution Bulletin, 76: 16–27.

444 Boyes, S. J., and Elliott, M. 2014. Marine legislation – The ultimate ‘horrendogram’: International law,
 445 European directives & national implementation. *Marine Pollution Bulletin*, 86: 39–47.

446 Cash, D.W., Clark, W.C., Alcock, F., Dickson, N.M., Eckley, N., Jäger, J., Mitchell, R.B., 2002. Saliency,
 447 Credibility, Legitimacy and Boundaries: Linking Research, Assessment and Decision Making. KSG Faculty
 448 Research Working Paper Series, RWP02-046. 25pp

449 Cisewski, B. and Strass, V. H. 2016. Acoustic insights into the zooplankton dynamics of the eastern Weddell
 450 Sea. *Progress in Oceanography*, 144: 42-92. doi:10.1016/j.pocean.2016.03.005.

451 Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R.V.,
 452 Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M. 1997. The value of the world's ecosystem services
 453 and natural capital. *Nature* 387:253-260

454 Cushman, S.A., Gutzweiler, K., Evans, J.S., McGarigal, K., 2010. The gradient paradigm: a conceptual and
 455 analytical framework for landscape ecology. In: Cushman, S.A. and Huettmann, F. (Eds) *Spatial*
 456 *complexity, informatics, and wildlife conservation*, Springer pp 83-108

457 Davidson, K., Miller, P., Wilding, T., Shutler, J., Bresnan, E., Kennington, K., and Swan, S. 2009. A large and
 458 prolonged bloom of *Karenia mikimotoi* in Scottish waters in 2006. *Harmful Algae*, 45: 692 – 703.

459 de Groot, R.S., Wilson, M.A., Boumans, R.M.J. 2002. A typology for the classification, description and
 460 valuation of ecosystem functions, goods and services. *Ecological Economics* 41: 393-408.

461 Della Penna, A., Koubbi, P., Cotté, C., Bon, C., Bost, C.-A., d'Ovidio, F. 2017. Lagrangian analysis of multi-
 462 satellite data in support of open ocean Marine Protected Area design. *Deep Sea Research Part II: Topical*
 463 *Studies in Oceanography* 140:212-221.

464 Dickey-Collas, M. 2014. Why the complex nature of integrated ecosystem assessments requires a flexible
 465 and adaptive approach. *ICES Journal of Marine Science*, 71: 1174–1182.

466 Etnoyer, P., Canny, D., Mate, B., and Morgan, L. 2004. Persistent pelagic habitats in the Baja California to
 467 Bering Sea (B2B) Ecoregion. *Oceanography*, 17: 1: 90-101.

468 European Commission, 2007. Communication from the commission to the European parliament, the council
 469 the European economic and social committee and the committee of the regions. An Integrated Maritime
 470 Policy for the European Union. Brussels COM(2007)575 final. 16pp

471 European Commission, 2008. Directive 2008/56/EC of the European Parliament and of the Council
 472 establishing a framework for community action in the field of marine environmental policy (Marine
 473 Strategy Framework Directive). *Official Journal of the European Union*, L164: 19–40.

474 European Commission, 2010. Commission Decision of 1 September 2010 on criteria and methodological
 475 standards on good environmental status of marine waters (notified under document C(2010)
 476 5956)(2010/477/EU). *Official Journal of the European Union*, L232: 12–24.

477 European Commission, 2017. Commission decision (EU) 2017/848 of 17 May 2017 laying down criteria and
 478 methodological standards on good environmental status of marine waters and specifications and
 479 standardised methods for monitoring and assessment, and repealing Decision 2010/477/EU. *Official*
 480 *Journal of the European Union*, L 125: 43-74.

481 Fernandes, J. A., Cheung, W. W. L., Jennings, S., Butenschön, M., de Mora, L., Frölicher, T. L., Barange, M.,
 482 and Grant, A. 2013. Modelling the effects of climate change on the distribution and production of marine

483 fishes: accounting for trophic interactions in a dynamic bioclimate envelope model. *Global Change*
484 *Biology*, 19: 2596-2607.

485 Funtowicz, S. and Ravetz, J. 1993. Science for the Post-Normal Age. *Futures* 25/7: 735-755.

486 Game, E. T., Grantham, H. S., Hobday, A. J., Pressey, R. L., Lombard, A. T., Beckley, L. E., Gjerde, K., *et al.*
487 2009. Pelagic protected areas: the missing dimension in ocean conservation. *Trends in Ecology &*
488 *Evolution*, 24: 360-369.

489 Gibbons, M., Limoges, C., Nowotny, H., Schwartzman, S., Scott, P., and Trow, M. 1994. The new production
490 of knowledge: the dynamics of science and research in contemporary societies. London: Sage. ISBN 0-
491 8039-7794-8.

492 Gillibrand, P. A., Siemering, B., Miller, P. I., and Davidson, K. 2016. Individual-based modelling of the
493 development and transport of a *Karenia mikimotoi* bloom on the North-west European continental shelf.
494 *Harmful Algae*, 53: 118 –134.

495 Gimpel, A., Stelzenmüller, V., Cormier, R., Floeter, J., and Temming, A. 2013. A spatially explicit risk approach
496 to support marine spatial planning in the German EEZ. *Marine Environmental Research*, 86: 56–69.

497 Gowen, R. J., Tett, P., Bresnan, E., Davidson, K., McKinney, A., Harrison, P. J., Milligan, S., Mills, D. K., Silke, J.,
498 and Crooks, A. M. 2012. Anthropogenic nutrient enrichment and blooms of harmful phytoplankton.
499 *Oceanography and Marine Biology: An Annual Review*, 50: 65–126.

500 Gray, J. S. 1997. Marine biodiversity: patterns, threats and conservation needs. *Biodiversity and*
501 *Conservation*, 6: 153-175.

502 Haury, L. R. 1976. A comparison of zooplankton patterns in the California Current and North Pacific Central
503 Gyre. *Marine Biology*, 37: 159-167.

504 HELCOM, 2010. Ecosystem Health of the Baltic Sea 2003–2007: HELCOM Initial Holistic Assessment. *Balt. Sea*
505 *Environ. Proc.* No. 122. 68pp.

506 Hidalgo, M., Reglero, P., Alvarez-Berastegui, D., Torres, A. P., Álvarez, I., Rodríguez, J. M., Carbonell, A., *et al.*
507 2015. Hidden persistence of salinity and productivity gradients shaping pelagic diversity in highly dynamic
508 marine ecosystems. *Marine Environmental Research*, 104: 47-50.

509 Hufnagl, M., Payne, M., Lacroix, G., Bolle, L.J., Daewel, U. *et al.* in press. Variation that can be expected when
510 using particle tracking models in connectivity studies. *Journal of Sea Research*.
511 doi.org/10.1016/j.seares.2017.04.009

512 Hyrenbach, K. D., Forney, K. A., and Dayton, P.K. 2000. Marine protected areas and ocean basin
513 management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 10: 437–458.

514 Jennings, S. 2005. Indicators to support an ecosystem approach to fisheries. *Fish and Fisheries* 6: 212–232.

515 Kavanaugh, M. T., Oliver, M. J., Chavez, F. P., Letelier, R. M., Muller-Karger, F. E., and Doney, S. C. 2016.
516 Seascapes as a new vernacular for pelagic ocean monitoring, management and conservation. *ICES Journal*
517 *of Marine Science*, 73: 1839–1850.

518 Knights, A. M., Koss, R. S., and Robinson, L. A. 2013. Identifying common pressure pathways from a complex
519 network of human activities to support ecosystem-based management. *Ecological Applications*, 23: 755–
520 765.

521 Kobayashi, D. R., Polovina, J. J., Parker, D. M., Kamezaki, N., Cheng, I. J., Uchida, I., Dutton, P. H., and Balazs,
522 G. H. 2008. Pelagic habitat characterization of loggerhead sea turtles, *Caretta caretta*, in the North Pacific
523 Ocean (1997–2006): Insights from satellite tag tracking and remotely sensed data. *Journal of*
524 *Experimental Marine Biology and Ecology*, 356: 96–114.

525 Liqueste, C., Piroddi, C., Drakou, E.G., Gurney, L., Katsanevakis, S., Charef, A., and Egoh, B. 2013. Current
526 Status and Future Prospects for the Assessment of Marine and Coastal Ecosystem Services: A Systematic
527 Review. *PLoS one* 8(7): e67737.

528 Louzao, M., Pinaud, D., Péron, C., Delord, K., Wiegand, T., and Weimerskirch, H. 2011. Conserving pelagic
529 habitats: seascape modelling of an oceanic top predator. *Journal of Applied Ecology*, 48: 121–132.

530 Mayor, S. J., Schaefer, J. A., Schneider, D. C., & Mahoney, S. P. 2007. Spectrum of selection: new approaches
531 to detecting the scale-dependent response to habitat. *Ecology*, 88(7), 1634-1640.

532 Manderson, J. P. 2016. Seascapes are not landscapes: an analysis performed using Bernhard Riemann’s rules.
533 *ICES Journal of Marine Science*, 73: 1831–1838.

534 McGarigal, K., Cushman, S.A. 2005. The gradient concept of landscape structure. In Wiens, J.A. and Moss,
535 M.R., (eds). *Issues and Perspectives in Landscape Ecology*. Cambridge University Press. pp 112-119

536 McQuatters-Gollop, A. 2012. Challenges for implementing the Marine Strategy Framework Directive in a
537 climate of macroecological change. *Philosophical Transactions of the Royal Society A*, 370: 5636–5655.

538 Mee, L., D., Jefferson, R. L., Laffoley, D. d’A., and Elliott, M. 2008. How good is good? Human values and
539 Europe’s proposed Marine Strategy Directive. *Marine Pollution Bulletin*, 56: 187–204.

540 Melián, C.J., Bascompte, J., and Jordano, P. 2005. Spatial structure and dynamics in a marine food web. pp.
541 19-24. In: Eds. A. Belgrano, U. M. Scharler, J. Dunne, and R. E. Ulanowicz. *Aquatic Food Webs - An*
542 *Ecosystem Approach*. Oxford University Press.

543 Möller, K. O., St. John, M. A., Temming, A., Flöter, J., Sell, A. F., Herrmann, J. P., and Möllmann, C. 2012.
544 Marine snow, zooplankton and thin layers: indications of a trophic link from small-scale sampling with the
545 Video Plankton Recorder. *Marine Ecology Progress Series*, 468: 57-69.

546 NOAA, 1996. National Oceanic and Atmospheric Administration Magnuson-Stevens Fishery Management
547 and Conservation Act amended through 11 October 1996. NOAA Technical Memorandum NMFS-F/SPO-
548 23.

549 OSPAR 2010. Quality Status Report 2010. OSPAR Commission London 176pp

550 Papathanasopoulou, E., Queirós, A. M., Beaumont, N., Hooper, T., and Nunes, J., 2016. What evidence exists
551 on the local impacts of energy systems on marine ecosystem services: a systematic map. *Environmental*
552 *Evidence*, 5:25 12pp. doi: 10.1186/s13750-016-0075-6.

553 Petitgas, P., Rijnsdorp, A.D., Dickey-Collas, M., Engelhard, G.H., Peck, M.A., Pinnegar, J.K., Drinkwater, K.,
554 Huret, M., Nash, R.D.M., 2013. Impacts of climate change on the complex life cycles of fish. *Fisheries*
555 *Oceanography* 22: 121–139.

556 Pittman, S. J., and Brown, K. A. 2011. Multi-scale approach for predicting fish species distributions across
557 coral reef seascapes. *PloS one*, 6(5), e20583.

558 Planque, B. 2015. Projecting the future state of marine ecosystems, “la grande illusion”? *ICES Journal of*
559 *Marine Science*, 73: 204-208.

560 Planque, B., Fromentin, J. -M., Cury, P., Drinkwater, K. F., Jennings, S., Perry, R. I., and Kifani, S. 2010. How
561 does fishing alter marine populations and ecosystems sensitivity to climate?, *Journal of Marine Systems*
562 79: 403-417, ISSN 0924-7963. <http://dx.doi.org/10.1016/j.jmarsys.2008.12.018>.

563 Purcell, E. M. 1977. Life at low Reynolds number. *American Journal of Physics*, 45, 3; doi: 10.1119/1.10903.

564 Queirós, A. M., Huebert, K. B., Key, F., Fernandes, J. A., Stolte, W., Maar, M., Kay, S., *et al.* 2016. Solutions for
565 ecosystem-level protection of ocean systems under climate change. *Global Change Biology*, 22: 3927–
566 3936, doi: 10.1111/gcb.13423.

567 Ray, G.C., 1991. Coastal-zone biodiversity patterns. *BioScience*, 41: 490-498.

568 Reiners, W.A., Lockwood, J.A., Reiners, D.S., and Prager, S.D., 2017. 100 years of ecology: what are our
569 concepts and are they useful? *Ecological Monographs*, 87: 260–277

570 Scales, K. L., Miller, P. I., Hawkes, L. A., Ingram, S. N., Sims, D. W., and Votier, S. C., 2014. On the front line:
571 frontal zones as priority at-sea conservation areas for mobile marine vertebrates. *Journal of Applied*
572 *Ecology*, 51: 1575–1583.

573 Scales, K.L., Hazen, E.L., Jacox, M.G., Edwards, C.A., Boustany, A.M., Oliver, M.J., and Bograd, S.J., 2017. Scale
574 of inference: on the sensitivity of habitat models for wide ranging marine predators to the resolution of
575 environmental data. *Ecography* 40: 210–220.

576 Schneider, D. C. 2001. The rise of the concept of scale in ecology. *Bioscience*, 51: 545-553.

577 Shephard, S., Rindorf, A., Dickey-Collas, M., Hintzen, N. T., Farnsworth, K., and Reid, D. G. 2014. Assessing the
578 state of pelagic fish communities within an ecosystem approach and the European marine strategy
579 framework directive. *ICES Journal of Marine Science*, 71: 1572–1585.

580 Shephard, S., Greenstreet, S. P. R., Piet, G. J., Rindorf, A., and Dickey-Collas, M. 2015. Surveillance indicators
581 and their use in implementation of the Marine Strategy Framework Directive. *ICES Journal of Marine*
582 *Science*, 72: 2269–2277.

583 Steele, J. H. 1978 *Spatial Pattern in Plankton Communities*. Plenum Press, New York, London.

584 Stommell, H. 1963. Varieties of oceanic experience. *Science* 139: 572-576.

585 Tett, P., Gowen, R. J., Painting, S. J., Elliott, M., Forster, R., Mills, D. K., Bresnan, E., *et al.* 2013. Framework for
586 understanding marine ecosystem health. *Marine Ecology Progress Series*, 494: 1–27.

587 Tew Kai, E., Rossi, V., Sudre, J., Weimerskirch, H., Lopez, C., Hernandez-Garcia, E., Marsac, F., Garçon, V.,
588 2009. Top marine predators track Lagrangian coherent structures. *Proceedings of the National Academy*
589 *of Sciences* 106:8245-8250

590 Treml, E. A., Roberts, J. J., Chao, Y., Halpin, P. N., Possingham, H. P., and Riginos, C. 2012. Reproductive
591 output and duration of the pelagic larval stage determine seascape-wide connectivity of marine
592 populations. *Integrative and Comparative Biology*, 52: 525–537.


593 Trenkel, V. M., Hintzen, N. T., Farnsworth, K. D., Olesen, C., Reid, D., Rindorf, A., Shephard, S., and Dickey-
594 Collas, M. 2015. Identifying marine pelagic ecosystem management objectives and indicators. *Marine*
595 *Policy*, 55: 23–32.

596 Trenkel, V. M., Handegard, N. O., and Weber, T. C. 2016. Observing the ocean interior in support of
597 integrated management. *ICES Journal of Marine Science*, 73: 1947–1954.

- 598 Turnhout, E., Hisschemöller, M., and Eijsackers, H. 2007. Ecological indicators: between the two fires of
599 science and policy. *Ecological Indicators*, 7, 215–228.
- 600 Vilchis, L. I., Balance, L. T., and Fiedler, P. C. 2006. Pelagic habitat of seabirds in the eastern tropical Pacific:
601 effects of foraging ecology on habitat selection. *Marine Ecology Progress Series*, 315: 279–292.
- 602 Wasmund, N., Kownacka, J., Göbel, J., Jaanus, A., Johansen, M., Jurgensone, I., Lehtinen, S., and Powilleit, M.
603 2017. The Diatom/Dinoflagellate index as indicator of ecosystem changes in the Baltic Sea 1. Principle and
604 handling instruction. *Frontiers in Marine Science*, 4: 13pp. doi: 10.3389/fmars.2017.00022.
- 605 Watson, J. R., Hays, C. G., Raimondi, P. T., Mitarai, S., Dong, C., McWilliams, J.C., Blanchette, C.A., *et al.* 2011.
606 Currents connecting communities: nearshore community similarity and ocean circulation. *Ecology*, 92:
607 1193–1200.
- 608 Young, K. V., Dower, J. F., and Pepin, P. 2009. A hierarchical analysis of the spatial distribution of larval fish
609 prey. *Journal of Plankton Research*, 31: 687-700.

610

611



612

613 Figure 1. Three key properties of the state of the pelagic system as contributors to GES.